

10/510444
PCT/GB/2003/000544

22 JUN 2005



INVESTOR IN PEOPLE



PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

RECEIVED	
10 FEB 2004	
WIPO	PCT

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated 9 January 2004

BEST AVAILABLE COPY

Patents Form 1/77

Patents Act 1977
(s 16)THE PATENT OFFICE
A
23 DEC 2002
RECEIVED BY FAXThe
Patent
OfficeEFFECTIVE
1/77**Request for grant of a patent***(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)*

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

IP/P7229

2. Patent application number

(The Patent Office will fill in this part)

0229734.9

3. Full name, address and postcode of the or of each applicant *(underline all surnames)*

QINETIQ LIMITED

Registered Office 85 Buckingham Gate
London SW1E 6PD
United KingdomPatents ADP number *(if you know it)*

8183857065

If the applicant is a corporate body, give the country/state of its incorporation

GB

4. Title of the invention

Grading Oestrogen and Progesterone Receptors Expression

5. Name of your agent *(if you have one)*

Bowdery Anthony Oliver

"Address for service" in the United Kingdom to which all correspondence should be sent *(including the postcode)*QINETIQ LIMITED
IP Formalities
A4 Bldg
Cody Technology Park
Ively Road
Farnborough
Hants GU14 0LX United KingdomPatents ADP number *(if you know it)*

8527376001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and *(if you know it)* the or each application number

Country

Priority application number
*(if you know it)*Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number or earlier application

Date of filing
*(day / month / year)*8. Is a statement of inventorship and of right to grant of a patent required in support of this request? *(Answer 'Yes' if:*

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.
See note (d))

YES (b)

Patents Form 1/77

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 15

Claim(s) 05

Abstract 01

Drawing(s) 02

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11. I / We request the grant of a patent on the basis of this application.

Signature

Mr P Davics

Date
23/12/02

12. Name and daytime telephone number of person to contact in the United Kingdom

Mrs Linda Bruckshaw 01252 392722

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent of the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have attached 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

Patents Form 1/77

Grading Oestrogen and Progesterone Receptors Expression

This invention relates to a method, an apparatus and a computer program for grading Oestrogen and Progesterone Receptors expression (ER and PR) in image data from histological slides to provide information for clinical treatment and prognosis. It is particularly (but not exclusively) relevant to providing clinical information on potentially cancerous tissue such as for example breast cancer tissue.

Breast cancer is a common form of female cancer. Once a lesion indicative of breast cancer has been detected, tissue samples are taken and examined by a histopathologist to establish a diagnosis, prognosis and treatment plan. However, pathological analysis of tissue samples is a time consuming and inaccurate process. It entails interpretation of images by human eye, which is highly subjective: it is characterised by considerable inaccuracies in observations of the same samples by different observers and even by the same observer at different times. For example, two different observers assessing the same ten tissue samples may easily give different opinions for three of the slides - 30% error. The problem is exacerbated by heterogeneity, i.e. complexity of some tissue sample features.

The hormone receptor status of breast cancer cells provides useful information for treatment and prognosis since the presence of some of these receptors can provide a means for controlling cell growth via chemotherapy. In general, cancers in which the cells express oestrogen receptor (ER) in their nuclei have a better prognosis and can respond to hormonal manipulation for example using the drug Tamoxifen. ER and PR are steroid hormone receptors. ER and PR expression can be observed in any tissue, i.e. tumour and non-tumour: they are used in breast cancer assessment because steroid hormone status guides selection of therapy. Almost 75% of breast cancers expressing ER will respond to this therapy, whereas less than 5% not expressing ER will respond. In general, cancers that are ER positive will also be PR positive. However, carcinomas that are PR positive, but not ER positive, may have a worse prognosis. Histological images produced for ER and PR determination are indistinguishable visually, though they are created by treating tissue samples with different chemical stains. Staining of tissue sample slides is obtained by using antibodies specific to respective receptors, i.e. one antibody for ER and a different

antibody for PR. Because ER and PR are both steroid receptors, their associated staining will be similar and so the same process can be used to extract both features.

There is a need to provide an objective measurement of ER and PR to support a pathologist's diagnosis and patient treatment.

- 5 The present invention provides a method of grading Oestrogen and Progesterone Receptors expression (ER and PR) from image data obtained from histological slides characterised in that it includes determining the number of relatively dark image pixels compared to relatively bright image pixels and providing a grading of ER or PR in accordance with the magnitude of the number of relatively dark pixels.
- 10 The invention provides the advantage that it is an objective procedure for providing a grading of ER or PR.

The number of relatively dark image pixels may be determined by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensities below a predetermined intensity threshold.

- 15 In another aspect, the present invention provides a method of grading ER and PR from image data obtained from histological slides characterised in that it includes the steps of:
- a) determining the number of pixels in an image having relatively dark intensities compared to other pixels in an image,
 - b) determining pixel number thresholds to quantify grading, and
 - 20 c) comparing the number of relatively dark pixels with the thresholds and providing a grading of ER or PR in accordance therewith.

In a further aspect, the present invention provides a method of grading ER and PR from image data obtained from histological slides characterised in that it includes the steps of:

- a) determining what proportion of total blob area is brown blob area in an image,
- 25 b) determining brown blob area proportion thresholds to quantify grading, and
- c) comparing the brown blob area proportion with the thresholds and providing a grading of ER or PR in accordance therewith.

The proportion of total blob area which is brown blob area may be determined by the steps of:

- 5 a) remapping pixel intensities in the image data to increase the contrast of relatively darker image regions and to transform relatively brighter image regions into a contrast-free background,
- b) converting the remapped image data into thresholded binary images from which total blob area and brown blob area are discernible respectively, and
- c) expressing brown blob area as a proportion of total blob area.

10 The step of providing a grading of ER or PR may provide a first contribution thereto and the method may include:

- a) providing a second contribution to grading of ER or PR by determining the number of relatively dark image pixels compared to relatively bright image pixels and deriving the second contribution in accordance with the magnitude of the number of relatively dark pixels, and
- 15 b) combining the first and second contributions.

The number of relatively dark image pixels may be determined by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensity below a predetermined intensity threshold.

20 In an alternate aspect, the present invention provides computer apparatus for grading ER and PR from image data obtained from histological slides characterised in that the apparatus is programmed to determine the number of relatively dark image pixels compared to relatively bright image pixels and to provide a grading of ER or PR in accordance with the magnitude of the number of relatively dark pixels.

25 The computer apparatus may be programmed to determine the number of relatively dark image pixels by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensities below a predetermined intensity threshold.

In a further alternative aspect, the present invention provides computer apparatus for grading ER and PR from image data obtained from histological slides characterised in that it is programmed to:

- 5 a) determine the number of pixels in an image having relatively dark intensities compared to other pixels in an image,
- b) determine pixel number thresholds to quantify grading, and
- c) compare the number of relatively dark pixels with the thresholds and provide a grading of ER or PR in accordance therewith.

10 In another alternative aspect, the present invention provides computer apparatus for grading ER and PR from image data obtained from histological slides characterised in that it is programmed to:

- a) determine what proportion of total blob area is brown blob area in an image,
- b) determine brown blob area proportion thresholds to quantify grading, and
- 15 c) compare the brown blob area proportion with the thresholds and provide a grading of ER or PR in accordance therewith.

The apparatus may be programmed to determine the proportion of total blob area which is brown blob area by:

- 20 a) remapping pixel intensities in the image data to increase the contrast of relatively darker image regions and to transform relatively brighter image regions into a contrast-free background,
- b) converting the remapped image data into thresholded binary images from which total blob area and brown blob area are discernible respectively, and
- c) expressing brown blob area as a proportion of total blob area.

25 The apparatus may be programmed to provide a grading of ER or PR as a first contribution thereto and also be programmed to:

- a) provide a second contribution to grading of ER or PR by determining the number of relatively dark image pixels compared to relatively bright image pixels and deriving the second contribution in accordance with the magnitude of the number of relatively dark pixels, and
- 30 b) combine the first and second contributions.

The apparatus may be programmed to determine the number of relatively dark image pixels by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensity below a predetermined intensity threshold.

- 5 In a different aspect, the present invention provides a computer programme for grading ER and PR from image data obtained from histological slides characterised in that it includes instructions for determining the number of relatively dark image pixels compared to relatively bright image pixels and providing a grading of ER or PR in accordance with the magnitude of the number of relatively dark pixels.
- 10 The programme may includes instructions for determining the number of relatively dark image pixels by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensities below a predetermined intensity threshold.

- In another different aspect, the present invention provides a computer programme for
- 15 grading ER and PR from image data obtained from histological slides characterised in that it includes instructions for:
- determining the number of pixels in an image having relatively dark intensities compared to other pixels in an image,
- determining pixel number thresholds to quantify grading, and
- 20 comparing the number of relatively dark pixels with the thresholds and providing a grading of ER or PR in accordance therewith.

- In a further different aspect, the present invention provides a computer programme for grading ER and PR from image data obtained from histological slides characterised in that it includes instructions for:
- 25 a) determining what proportion of total blob area is brown blob area in an image,
- b) determining brown blob area proportion thresholds to quantify grading, and
- c) comparing the brown blob area proportion with the thresholds and providing a grading of ER or PR in accordance therewith.

The programme may include instructions for determining the proportion of total blob area which is brown blob area by:

- remapping pixel intensities in the image data to increase the contrast of relatively darker image regions and to transform relatively brighter image regions into a contrast-free background,
- 5 converting the remapped image data into thresholded binary images from which total blob area and brown blob area are discernible respectively, and
- expressing brown blob area as a proportion of total blob area.

The instructions for providing a grading of ER or PR may be for providing a first contribution thereto and the programme may also include instructions for:

- 10 a) providing a second contribution to grading of ER or PR by determining the number of relatively dark image pixels compared to relatively bright image pixels and deriving the second contribution in accordance with the magnitude of the number of relatively dark pixels, and
- 15 b) combining the first and second contributions.

The programme may include instructions for determining the number of relatively dark image pixels by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensity below a predetermined intensity threshold.

20 In order that the invention might be more fully understood, embodiments thereof will now be described, by way of example only, with reference to the accompanying drawings, in which:-

Figure 1 is a block diagram of a procedure of the invention for grading oestrogen and progesterone receptors expression; and

25 Figure 2 is a block diagram showing part of the procedure of Figure 1 in more detail.

Referring to Figure 1, there is illustrated a procedure 10 of the invention for grading oestrogen (ER) and progesterone (PR) receptors expression in tissue samples presented as histopathological slides of potential carcinomas of the breast. The procedure 10 requires

data from histological slides in a suitable form. Sections are taken (cut) from breast tissue samples (biopsies) and placed on respective slides. Slides are stained using the ER staining antibody-peroxidase (P/ER)- to assess oestrogen receptor status, which is a stain used to indicate uptake of oestrogen. In the present example, image data were obtained by a pathologist using Zeiss Axioskop microscope with a J0enoptiks Progres 3012 digital camera. Image data from a slide is a set of digital images obtained at a linear magnification of 40 (i.e. 40X).

To select images, a pathologist scans the microscope over a slide, and at 40X magnification selects a region (referred to as a tile) of the slide which appears to be most promising in terms of an analysis to be performed. This region is then photographed using the microscope and digital camera referred to above. The digital camera produces a digitised image in three colours, i.e. red, green and blue (R, G & B) intensity values for each pixel each in eight bits and therefore in the range 0 to 255: each image is an electronic equivalent of a tile. Three intensity values are obtained for each pixel in a pixel array to provide a colour image as a combination of R, G and B image planes. The image data from the tile is stored in a database 12 for later use. Oestrogen (ER) and progesterone (PR) receptors expression are determined using a feature detection process 14: this provides a score 18 for input to a diagnostic report at 20.

ER and PR are measured from brown stain characteristics (blob colour intensity and area ratio) in a specimen produced using tissue staining processes mentioned above. The objective of the procedure 10 is to quantify cancer cells' response to oestrogen and progesterone: this uses the fact that the more oestrogen or progesterone positive a specimen is the greater the intensity and area of its brown stain.

In a prior art manual procedure, a clinician places a slide under a microscope and examines a region of it at magnification of x40 for indications of ER and PR. The prior art manual procedure for scoring ER and PR involves a pathologist subjectively estimating the stained surface area proportion as well as staining intensity. The process described below in this example replaces the prior art manual procedure with an objective procedure.

Referring now to Figure 2, the process 14 is shown in more detail. At 30 an input red, green and blue (RGB) digital colour image (initial image designated I) is a starting point for obtaining a score for stained area proportion. The image I has three eight-bit values in the range 0 to 255 per pixel for the R, G and B image planes. In this image, cells in a tissue sample appear in a variety of colours, shapes and sizes, and are referred to as "blobs" to reflect this. Blobs appearing brown are of interest for ER/PR measurement. The "stained area proportion" measurement is the sum of the areas of brown stained blobs divided by the sum of the areas of all blobs in the image.

It has been found that pixels of interest for the purposes of assessing staining are those with relatively low intensity values. The next stage 32 is therefore to transform the initial image I so that an intensity range 0-100 of its darker pixels in each of the three R, G and B image planes is transformed by a mapping to fill the full eight-bit intensity range 0 to 255; this means that a pixel intensity value of 100, which is less than half but more than one quarter of the 255 maximum intensity value, becomes transformed to 255. It also has the effect of transforming pixels in I with intensity greater than 100 to 255. If a pixel has R, G and B all equal to 255, it appears white in the image.

In this example, specimen images were processed using computer software referred to as "Matlab[®]" produced by Mathworks Inc., an American corporation. This software provides a variety of image processing functions. Blobs are identified in the initial image I using the Matlab "imadjust" function, which maps an image's intensity values to a new range. Imadjust is used as previously indicated to remap darker pixel intensity values of all three I image planes R, G and B in the range 0 to 100 to fill the full intensity range of 0 to 255; this increases the contrast of darker regions of the image as well as transforming into a contrast-free white background those pixels with relatively high intensities of 100 and above in all image planes. This transformation is shown in Table 1 below.

Table 1: Image transformation

Input image I: R, G or B pixel value P_I	Output image K: R, G or B pixel value P_K
$0 \leq P_I \leq 100$	$P_K = 255P_I/100$
$P_I \geq 100$	$P_K = 255$

The syntax of `imadjust` for mapping the input image *I* to an output image *K* is:

$$K = \text{imadjust}(I, [\text{low_in } \text{high_in}], [\text{low_out } \text{high_out}]) \quad (1)$$

- 5 where `low_in` and `high_in` are the chosen dark intensities 0 and 100 in the input image *I* which are mapped to `low_out` and `high_out` 0 and 255 in the output image *K* at 32. Because only relatively darker pixels are retained, it is assumed that they approximately correspond to all blobs whether stained or not. A bitmap image is consequently obtained at 34 using another Matlab function referred to as "`im2bw`", which creates from the RGB image *K* a
10 binary bitmap image denoted *L* with pixel values 0 and 1 derived using a luminance threshold of 0.9.

- To derive *L*, the image *K* is firstly converted from red/green/blue (RGB) to a one colour image. In this example the conversion of *K* is by transformation into a different image space hue/saturation/value (HSV). Value is also referred to as brightness, luminance or
15 intensity. The RGB to HSV transformation is well known and described in more detail on page 77 of *Image Based Measurement Systems*, F. Van Der Heijden. It will not be described further. It is not essential to use HSV, one could select one of the image planes or use another known transformation such as principal component analysis (see e.g. Jackson J.E., 'A User Guide to Principal Components' pp 1-25, John Wiley & Sons, 1991).
20 In the present example a one colour image is derived by setting to zero the hue *H* and saturation *S* components of pixels in the transformation of *K*, and the value (luminance) component *V* is retained as a greyscales image. *V* has values in the range 0 to 1.0. All pixels with a value greater than or equal to 0.9 are then set to 1 and those less than 0.9 are set to 0: this creates the binary bitmap image *L* in which pixels of blobs are 0 and
25 background pixels are 1.

- The sum of the areas of all blobs in image *I* collectively (i.e. whether stained brown or not) is denoted by *S1*. *S1* is computed by inverting the bitmap image *L* (exchanging 1 for 0 pixel values and vice versa) to form an image L^{-1} (or "`not(L)`" in Matlab) in which blob pixels are value 1; pixels in L^{-1} which are not blob pixels are 0. The value 1 pixels in L^{-1}
30 are then counted at 34 to provide *S1*, which is given in Matlab by the instruction:

$$\text{i.e. } S1 = \text{sum}(\text{sum}(\text{not}(L))) \quad (2)$$

The sum of the areas of all blobs stained brown in image I denoted S2 is now required. Here again the image K (see at (1) above) is converted at 36 to a single colour greyscales image, but for this purpose the blue image plane of the image K is selected: this is because trials showed that the blue image plane contains more brown staining information and fewer spurious details than the green and red planes. Using the im2bw Matlab function, as before the single colour image pixel values (in this case the blue image plane) are converted from the range 0 to 255 to 0 to 1 by division by the maximum of the pixel values, normally 255. A second binary bitmap image M is obtained from the 0 to 1 pixel value equivalent of the single colour image by thresholding: i.e. all pixels with a value greater than or equal to a threshold of 0.85 are set to 1 and those less than 0.85 are set to 0. This threshold provides for removal (merging into background) of blob pixels originally bluish and retention of those originally brownish. The sum S2 of the areas of all blobs stained brown is computed at 36 by inverting the bitmap image M to form image M⁻¹ (or "not(M)"), and counting the pixels which are equal to 1 in M⁻¹ and which are therefore blob pixels; here again pixels in M⁻¹ which are not brownish blob pixels are 0. S2 is given by the instruction:

$$S2 = \text{sum}(\text{sum}(\text{not}(M)))$$

(3)

The percentage W of blobs or cells stained brown is then computed at 38 by dividing S2 by S1 and multiplying by 100:

$$W = 100S2/S1$$

(4)

The score for proportion staining is denoted as SC1 and is obtained at 38 by comparing W to the thresholds mentioned above. For this example, the score for proportion of brown staining of cells in ER and PR detection in breast cancer is tabulated in Table 2 below.

Table 2: Score for proportion of brown staining:

W: % of cells stained brown	Proportion Score SC1
0 - 0.3%	0
0.3% < W < 1%	1
1-10%	2
11-33%	3
34-66%	4
67-100%	5

The thresholds in Table 2 are largely as used in manual assessment, except that the thresholds for scores 0 and 1 have been changed slightly from the manual equivalents (0% efficient than the manual equivalents using a specific set of trial images in testing the invention.

The next stage 40 is to obtain a score SC2 for cell staining intensity. A study of a set of trial images with different staining intensity for the purposes of the invention has shown that the occurrence frequency of relatively darker pixels (intensity value ≤ 100) in the RGB initial image I varies sufficiently with intensity staining to provide a measurement technique in this regard. In particular, strong staining is characterised by a relatively high occurrence frequency of darker pixels, but this frequency is zero (no darker pixels) when there is no staining. For weak to moderate staining, the frequency is intermediate. Consequently, in accordance with the invention, the next step is to form a histogram showing pixel intensity occurrence frequency: to achieve this, the RGB initial image I undergoes an HSV transformation of the kind described earlier. Pixel intensities in the V (value or intensity) image plane of the transformed image are then analysed to produce the histogram. The histogram has an abscissa axis with 256 pixel intensity values or bins and an ordinate axis showing number of pixels with the intensity associated with the respective bin each case. Based on these observations, it was decided to sum the numbers of pixels

contained in a sufficient number of the lower valued (darker) pixel intensity bins to give a statistically significant result. The number of bins is not very critical and in this example the bins that were chosen were bins 1 to 100 inclusive: this corresponds to counting pixels with intensities below a intensity threshold equal to the intensity associated with bin number 101. Here again Matlab functions were used such as "imhist" that returns the histogram as bin counts and bin locations as x (ordinate) values. A Matlab routine was written to carry out the summation of pixels in bins 1 to 100.

The score SC2 for intensity staining was derived using thresholds derived from analysis of a set of images and partly from the value of SC1. Denoting the total number of pixels in bins 1 to 100 collectively as NP(1:100), SC2 was 3 if NP(1:100) was greater than 100,000. SC2 was 2 if NP(1:100) was in the range 2001 to 100,000, and it was 1 if the number of pixels was found to be in the range 400 and 2000. If the number of pixels was less than 400, SC2 would be 0 if SC1 was 4 or less and 1 if SC1 was 5. This is tabulated below in Table 3.

Table 2: Score for intensity of staining:

NP(1:100): Number of Pixels in Bins 1 to 100	SC1	SC2	Staining
> 100,000	Any	3	Strong
2,001 to 100,000	Any	2	Moderate
400 to 2,000	Any	1	Weak
< 400	5	0	Weak
< 400	0 to 4	0	None

Having obtained SC1 and SC2, they are summed at 42 to provide an overall intermediate score in the range 0 to 8. The overall score is subject to the condition that if SC1 or SC2 is

0, the overall score is also set to 0. This is because a zero percentage of brown blobs implies zero staining intensity and vice versa. In principle one could obtain an estimate of ER or PR using SC1 or SC2 alone, but results are better when SC2 is combined with SC1 as described above.

- 5 In ER and PR determination there is a convention that an overall final score be determined in the range 0 to 3 from the overall intermediate score as set out in Table 4 below.

Overall Intermediate Score	Overall Final Score	Endocrine Treatment Prospects
0	0	Failure to respond
2 or 3	1	Small Chance of Response
4, 5 or 6	2	50% Chance of Response
7 or 8	3	High Chance of Response

The procedure of the invention is rapid. One image takes on average 9 seconds to process. It has been evaluated for an overall number of 97 images for ER and 92 images for PR.

- 10 Results produced by use of the invention with these images have been compared to assessments made manually by a pathologist, and are very encouraging. Over 80% of scores obtained as described above agreed with a pathologist's overall final score, and over 90% either agreed with the pathologist's score or differed by not more than 1.

- 15 It has been found that a scoring failure may result from confusion between slides stained very dark blue and those stained very brown. To counteract this, a correction for the overall final score has been developed based on image coloration. The correction involves transforming the RGB initial image I to a hue, saturation, value (HSV) image of the kind described earlier. Colour perception usually refers to three quantities that are hue, saturation and brightness: hue refers more specifically to a dominant wavelength or colour
- 20 in an image and thus it is of interest for discriminating between blue and brown.

Hue provides a check on whether blobs treated as being brown are indeed brown, or instead dark blue as sometimes happens with badly stained images. A parameter called "rep" is obtained by summing the values of pixels of blobs thought to be brown. Blobs thought to be brown are indicated by pixels with value 1 in the thresholded binary M^{-1} (in
5 Matlab code "not(M)") previously defined. A hue or H image plane H of I is derived at 44 for the initial image I by HSV transformation as aforesaid and rejection of the S and V values. Each pixel value in H of I is then multiplied by a respective pixel value in the same location in M^{-1} to yield HB of I. All non-blob pixels in HB of I are 0 and all blob pixels have respective values as in H of I. The required parameter rep is now obtained at 44 once
10 more by summing the pixel values in HB of I: this sum - i.e. rep - was observed to vary considerably depending on whether blobs have been stained predominantly blue or predominantly brown. A small value of rep is usually characteristic of bluish slides while a large value of rep is characteristic of brownish slides. Rep therefore provides a test of blueness or brownness enabling a correction to be applied to an overall intermediate score
15 in the range 0 to 8: the correction increases or decreases the overall intermediate score according to what rep indicates the associated slide staining colour to be.

For example, a small value of rep such as 0.05 indicates bluish images not containing many brown pixels. An overall intermediate score of 7 for an image associated with a small value of rep is corrected at 46 by being decreased by 1 or 2 to reflect the relative
20 paucity of brown pixels which its rep value indicates: such an image is unlikely to merit an overall intermediate score of 7 (overall final score 3) but is more likely to merit an overall intermediate score of 6 or 5 (overall corrected final score 2). It has been found that this correction improves results for both ER and PR. For PR in particular, 88% of corrected overall final scores agreed with the pathologist's manual result, and none differed from the
25 latter by more than 1. Since pathologist's manual results can be up to 30% in error as mentioned earlier, the difference may not be statistically significant. However, these results were obtained using a limited data set. It is desirable to verify the procedure of the invention using more images, laboratories and staining procedures.

The example of the invention in the foregoing description can clearly be implemented by
30 an appropriate computer program comprising program instructions recorded on an appropriate carrier medium and running on a conventional computer system. As described,

much of the program requirements are commercially available in Matlab, and can easily be augmented. The carrier medium may be a memory, a floppy or compact or optical disc or other hardware recordal medium, or an electrical signal. Such a program is straightforward for a skilled programmer to implement without requiring invention, because it involves
5 well known computational procedures.

CLAIMS

1. A method of grading Oestrogen and Progesterone Receptors expression (ER and PR) from image data obtained from histological slides characterised in that it includes determining the number of relatively dark image pixels compared to relatively bright image pixels and providing a grading of ER or PR in accordance with the magnitude of the number of relatively dark pixels.
2. A method according to Claim 1 characterised in that the number of relatively dark image pixels is determined by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensities below a predetermined intensity threshold.
3. A method of grading ER and PR from image data obtained from histological slides characterised in that it includes the steps of :
 - a) determining the number of pixels in an image having relatively dark intensities compared to other pixels in an image,
 - b) determining pixel number thresholds to quantify grading, and
 - c) comparing the number of relatively dark pixels with the thresholds and providing a grading of ER or PR in accordance therewith.
4. A method of grading ER and PR from image data obtained from histological slides characterised in that it includes the steps of:
 - a) determining what proportion of total blob area is brown blob area in an image,
 - b) determining brown blob area proportion thresholds to quantify grading, and
 - c) comparing the brown blob area proportion with the thresholds and providing a grading of ER or PR in accordance therewith.
5. A method according to Claim 4 characterised in that the proportion of total blob area which is brown blob area is determined by the steps of:

- a) remapping pixel intensities in the image data to increase the contrast of relatively darker image regions and to transform relatively brighter image regions into a contrast-free background,
 - b) converting the remapped image data into thresholded binary images from which total blob area and brown blob area are discernible respectively, and
 - c) expressing brown blob area as a proportion of total blob area.
6. A method according to Claim 3, 4 or 5 characterised in that the step of providing a grading of ER or PR provides a first contribution thereto and the method includes:
 - a) providing a second contribution to grading of ER or PR by determining the number of relatively dark image pixels compared to relatively bright image pixels and deriving the second contribution in accordance with the magnitude of the number of relatively dark pixels, and
 - b) combining the first and second contributions.
7. A method according to Claim 6 characterised in that the number of relatively dark image pixels is determined by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensity below a predetermined intensity threshold.
8. Computer apparatus for grading ER and PR from image data obtained from histological slides characterised in that the apparatus is programmed to determine the number of relatively dark image pixels compared to relatively bright image pixels and to provide a grading of ER or PR in accordance with the magnitude of the number of relatively dark pixels.
9. Apparatus according to Claim 8 characterised in that it is programmed to determine the number of relatively dark image pixels by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensities below a predetermined intensity threshold.
10. Apparatus for grading ER and PR from image data obtained from histological slides characterised in that it is programmed to:

- a) determine the number of pixels in an image having relatively dark intensities compared to other pixels in an image,
 - b) determine pixel number thresholds to quantify grading, and
 - c) compare the number of relatively dark pixels with the thresholds and provide a grading of ER or PR in accordance therewith.
11. Apparatus for grading ER and PR from image data obtained from histological slides characterised in that it is programmed to:
- a) determine what proportion of total blob area is brown blob area in an image,
 - b) determine brown blob area proportion thresholds to quantify grading, and
 - c) compare the brown blob area proportion with the thresholds and provide a grading of ER or PR in accordance therewith.
12. Apparatus according to Claim 11 characterised in that it is programmed to determine the proportion of total blob area which is brown blob area by:
- a) remapping pixel intensities in the image data to increase the contrast of relatively darker image regions and to transform relatively brighter image regions into a contrast-free background,
 - b) converting the remapped image data into thresholded binary images from which total blob area and brown blob area are discernible respectively, and
 - c) expressing brown blob area as a proportion of total blob area.
13. Apparatus according to Claim 10, 11 or 12 characterised in that it is programmed to provide a grading of ER or PR as a first contribution thereto it is also programmed to:
- a) provide a second contribution to grading of ER or PR by determining the number of relatively dark image pixels compared to relatively bright image pixels and deriving the second contribution in accordance with the magnitude of the number of relatively dark pixels, and
 - b) combine the first and second contributions.
14. Apparatus according to Claim 13 characterised in that it is programmed to determine the number of relatively dark image pixels by transforming the image

data to a different image space having an intensity image plane and counting the number of pixels having intensity below a predetermined intensity threshold.

15. A computer programme for grading ER and PR from image data obtained from histological slides characterised in that it includes instructions for determining the number of relatively dark image pixels compared to relatively bright image pixels and providing a grading of ER or PR in accordance with the magnitude of the number of relatively dark pixels.
16. A programme according to Claim 15 characterised in that it includes instructions for determining the number of relatively dark image pixels by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensities below a predetermined intensity threshold.
17. A computer programme for grading ER and PR from image data obtained from histological slides characterised in that it includes instructions for:
 - a) determining the number of pixels in an image having relatively dark intensities compared to other pixels in an image,
 - b) determining pixel number thresholds to quantify grading, and
 - c) comparing the number of relatively dark pixels with the thresholds and providing a grading of ER or PR in accordance therewith.
18. A computer programme for grading ER and PR from image data obtained from histological slides characterised in that it includes instructions for:
 - a) determining what proportion of total blob area is brown blob area in an image,
 - b) determining brown blob area proportion thresholds to quantify grading, and
 - c) comparing the brown blob area proportion with the thresholds and providing a grading of ER or PR in accordance therewith.
19. A programme according to Claim 18 characterised in that it includes instructions for determining the proportion of total blob area which is brown blob area by:

- a) remapping pixel intensities in the image data to increase the contrast of relatively darker image regions and to transform relatively brighter image regions into a contrast-free background,
 - b) converting the remapped image data into thresholded binary images from which total blob area and brown blob area are discernible respectively, and
 - c) expressing brown blob area as a proportion of total blob area.
20. A programme according to Claim 17, 18 or 19 characterised in that the instructions for providing a grading of ER or PR are for providing a first contribution thereto and it also includes instructions for:
- a) providing a second contribution to grading of ER or PR by determining the number of relatively dark image pixels compared to relatively bright image pixels and deriving the second contribution in accordance with the magnitude of the number of relatively dark pixels, and
 - b) combining the first and second contributions.
21. A programme according to Claim 20 characterised in that it includes instructions for determining the number of relatively dark image pixels by transforming the image data to a different image space having an intensity image plane and counting the number of pixels having intensity below a predetermined intensity threshold.

ABSTRACT

A method of grading Oestrogen and Progesterone Receptors expression (ER and PR) from histological images determines the percentage of brown image blob area in total blob area and derives percentage thresholds to quantify grading. Brown blob area proportion is then compared with the thresholds to provide a first contribution to grading of ER or PR. The number of relatively dark pixels is then counted, pixel number thresholds are determined to quantify grading, and the number of relatively dark pixels is compared with the thresholds to provide a second contribution to grading of ER or PR. The two contributions are then added to provide an overall grading, which is then scaled to the conventional range 0 to 3.

10

RECEIVED 13 OCT 1991





Figure 1

1/2



14 →

Figure 2

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.